




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification⁵ : A23J 1/12, 3/14, C07K 3/02 C07K 3/12, 3/28, 15/10	A1	(11) International Publication Number: WO 93/12667 (43) International Publication Date: 8 July 1993 (08.07.93)
(21) International Application Number: PCT/US92/11134 (22) International Filing Date: 18 December 1992 (18.12.92) (30) Priority data: 814,118 26 December 1991 (26.12.91) US (71) Applicant: OPTA FOOD INGREDIENTS, INC. [US/ US]; 64 Sidney Street, Cambridge, MA 02139 (US). (72) Inventors: COOK, Richard, B. ; 7 Agawam, Revere, MA 02151 (US). MALLEE, Frank, M. ; 3 Lisa Lane, Acton, MA 01720 (US). SHULMAN, Mark, L. ; 79 Central Street, No. 1, Waltham, MA 02154 (US).		(74) Agents: CARROLL, Alice, O. et al.; Hamilton, Brook, Smith & Reynolds, Two Militia Drive, Lexington, MA 02173 (US). (81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>With amended claims.</i>

(54) Title: PURIFICATION OF ZEIN FROM CORN GLUTEN MEAL**(57) Abstract**

Zein, zein bodies, glutelins or destarched corn gluten are extracted and purified from corn gluten meal by efficiently removing and recovering food grade pigments and starch hydrolysates, while discarding unwanted fractions. Corn gluten is subjected to the combination of enzymatic starch hydrolysis, alkaline treatment, alcohol washing, alcohol extraction, carbon or ion exchange resin purification, and precipitation to produce a substantially starch-free, deflavored and decolorized zein which is suitable for use in foods and pharmaceuticals. An advantage of the process is its use of food grade alcohol(s) to deflavor, decolorize, and extract the zein from corn gluten meal. Deflavored, decolorized and destarched corn gluten, zein bodies or glutelins are also obtained and are useful as vegetable protein supplements. By-products of the invention such as yellow food pigments and starch hydrolysates can be further purified and/or used directly in various foods.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NL	Netherlands
BE	Belgium	GN	Guinea	NO	Norway
BF	Burkina Faso	GR	Greece	NZ	New Zealand
BG	Bulgaria	HU	Hungary	PL	Poland
BJ	Benin	IE	Ireland	PT	Portugal
BR	Brazil	IT	Italy	RO	Romania
CA	Canada	JP	Japan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SK	Slovak Republic
CI	Côte d'Ivoire	LJ	Liechtenstein	SN	Senegal
CM	Cameroon	LK	Sri Lanka	SU	Soviet Union
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	MC	Monaco	TG	Togo
DE	Germany	MG	Madagascar	UA	Ukraine
DK	Denmark	ML	Mali	US	United States of America
ES	Spain	MN	Mongolia	VN	Viet Nam
FI	Finland				

-1-

PURIFICATION OF ZEIN FROM CORN GLUTEN MEALBackground of the Invention

A primary by-product of commercial corn wet milling is wet corn gluten meal (CGM) which is typically concentrated and sold in dry form (approximately 90% solids). CGM is known to be most stable in dry form. Zein can be readily obtained by further processing of corn gluten meal. The potential yield of zein is about one pound per bushel of corn.

Corn gluten has been processed into varying grades of zein depending upon the particular end use, i.e., for use in plastics, protective coatings, inks, food products, pigments, adhesives and films, among other uses. In food applications, the quality of the zein is of paramount importance to the overall organoleptic properties of the food product. For instance, currently available grades of zein contain impurities that can impart off-flavors and colors (particularly yellow color - due to the presence of xanthophyll and carotene pigments) to the food product. Further processing of the zein to remove undesirable colors and off-flavors has been developed with limited success, particularly for use in foods.

Processes for producing "white" zein by extraction with ethylene dichloride (Mason and Palmer, J. Biol. Chem. 107:131-132 (1934)), benzene (U.S. Patent No. 2,133,591 issued to Swallen) and other hydrocarbons (U.S. Patent Nos. 2,287,649 and 2,272,488 both issued to Swallen; U.S. Patent No. 2,229,870 issued to Pearce) have been described. Although the product has minimal color, the use of these solvents make the "white" zein unacceptable for use in the food and pharmaceutical industry.

-2-

Enzymatic modification of corn gluten by proteolytic hydrolysis to produce a flour-like material was previously described by Adler-Nissen (U.S. Patent 4,100,151). Enzyme hydrolysis of crude gluten using alpha-amylase has also been described by Knight (U.S. Patent No. 3,782,964). In both of these processes, the product generated is a modified gluten rather than zein.

Alcohol extraction of corn gluten to obtain zein has been described by Lawhorn (U.S. Patent No. 4,624,805), Chen and Hoff (U.S. Patent No. 4,716,218), Coleman (U.S. Patent No. 2,355,056), Carter and Reck (U.S. Patent No. 3,535,305) and Swallen (U.S. Patent Nos. 2,287,649, 2,332,356, 2,105,760, 2,120,946, 2,133,591 and 2,272,488). In each instance, zein is extracted using a solvent (e.g., ethanol, isopropanol) and concentration for which zein is soluble therein. The extracted zein produced by these methods tend to contain impurities that are co-extracted and soluble in the extraction solvent. Further extraction of the impurities is necessary to adequately depigment and de Flavor the zein product.

Thus, it is desirable to process corn gluten to generate a purified, food grade zein that is essentially decolorized, de Flavored and odor-free and suitable for use in food and pharmaceutical applications.

Summary of the Invention

This invention pertains to methods for purifying, fractionating and recovering zein, zein bodies, glutelins and mixtures thereof from corn gluten meal. Purification involves removing and recovering food grade pigments, glutelin(s) and starch hydrolysates, while discarding unwanted fractions. Corn gluten is subjected to a

-3-

combination of enzymatic starch hydrolysis, alkaline treatment, alcohol washing and either deflavored, decolored alcohol extraction of the deflavored and decolored zein or fractionation of the resultant deflavored, decolored gluten into zein body or glutelin components. The deflavored and decolored zein is suitable for use in foods and pharmaceuticals. An advantage of the process described herein is its use of food grade alcohol(s) to deflavor, decolor and extract the zein from corn gluten meal. Purified zein and zein bodies have improved utility and functionality in a variety of existing and emerging food applications. By-products such as glutelins, pigments and starch hydrolysates are also obtained by the methods of this invention and can be further purified for use as natural food colors and vegetable protein supplements. Alternatively, zein bodies may be harvested separate from glutelins or admixed with them in various proportions as deflavored and decolored corn gluten.

Brief Description of the Figures

Figure 1 is a schematic representation of the zein purification process of this invention.

Figure 2 is a graphic representation of cloud point temperatures for 30% zein solutions in varying concentrations of aqueous alcohol (e.g., ethanol, methanol, isopropanol and n-butanol). R.A. Reiners, et al., "Corn Proteins: Potential For Their Industrial Use", 58th American AOCC Meeting, St. Louis, MO, November 4, 1973.

Figure 3 shows the results from a high titer ethanol depigmentation wash. Corn gluten meal was washed with 100% Ethanol at 23°C (□) and 45°C (o). In the sixth extraction, water was used instead of ethanol.

-4-

Figure 4 shows two high performance liquid chromatograms obtained on aqueous extracts of commercial zein (Freeman Industries, Inc., Tuckahoe, New York) and zein produced by the methods of this invention (solid line just above x-axis). Commercial zein had 18 peaks with a peak area of 149.79×10^6 units. The purified zein had zero peaks and a peak area of zero units.

Detailed Description of the Invention

This invention pertains to methods for purifying zein from corn gluten meal, a major by-product of wet corn milling. Either a wet or gently dried gluten can be used as the starting material for the methods of this invention. The term "wet corn gluten" refers to an aqueous mixture of corn gluten that is muddy in appearance and contains from about 10-50% solids (e.g., from about 50 to about 75% protein and from about 15 to about 30% starch). Wet corn gluten also contains an amount of sulfite (approx. over 1,000 ppm dry weight) which is beyond the maximum permissible level for most foods, (100ppm limit for shrimp, for example). The term "dry corn gluten" refers to a corn milling by-product that has been further processed to reduce its moisture content to stabilize the corn gluten against degradation. This form of gluten, however, should be gently dried under conditions which preserve zein body membranes and spheroidal shape. Typically dry corn gluten (also referred to as "commercial corn gluten") contains about 90% solids. Wet or gently dried forms of corn gluten that deviate from the above description are included within the scope of this invention. The wet form of gluten is preferred for the purpose of this invention because the expense of drying and the generation of off-flavors associated with

-5-

high temperature (90-400°C) conventional drying are eliminated. However, the quality of wet gluten for the purpose of this invention, will deteriorate after a few weeks if it is not stored frozen. In comparison, commercial (or dry) CGM which has heretofore been the preferred raw material for obtaining commercial zein, offers the advantage of improved microbial stability and shelf life at ambient temperature.

According to this invention, corn gluten meal (either wet or gently dried forms) is subjected to a series of treatments which include enzymatic digestion of starch, alkaline removal of hydrocarbons and off-flavors, separate pigment and zein extraction with alcohol, leaving glutelin, alcohol washes, extractions and centrifugal separation to remove pigments, glutelin(s), residual starch and hydrocarbons, to thereby yield deflavored and decolored zein. These contaminants contribute colors, off-flavors and odors to zein. In the broadest aspect of the invention, deflavored and decolored zein or zein bodies and glutelins are produced from corn gluten by first removing color and flavor impurities from the gluten using an aqueous alcoholic solvent having a concentration which does not substantially extract the zein, zein bodies and glutelin therefrom. The alcohol concentration is then changed to separate the gluten into deflavored and decolored zein or zein bodies and glutelin fractions. Figure 1 schematically illustrates the steps employed to purify zein from wet or gently dried corn gluten. Each of these steps are specifically described in detail below.

Wet or gently dried corn gluten, obtained either from a commercial source or directly from a wet milling product stream, is first subjected to enzymatic treatment

-6-

to hydrolyze residual starch in the corn gluten. Complete digestion and removal of starch from the gluten substantially improves the effectiveness of the subsequent decoloring and deflavoring steps due, in part, to proximity of starch granules, zein bodies and glutelins in the corn gluten. Enzymatic hydrolysis is accomplished primarily through the action of α -amylase. In addition to α -amylase, this step can be performed using a combination of other carbohydrate hydrolyzing enzymes, such as glucoamylase, pullulanase or cellulase. The destarching step can be optimized by first homogenizing or agitating the corn gluten meal suspension to adequately separate and disperse the particles. Brief homogenization should be employed when wet CGM is used. Homogenization, however, is especially necessary when dry corn gluten is used as the starting material. Care should be taken to avoid harsh homogenization conditions (e.g. high pressure homogenizers) that can rupture the native structure of cell-like or sub-cellular organelles, such as zein body membranes.

By-product starch hydrolysates can be recovered from the discarded portion and further processed and/or purified. The yield of carbohydrate by-product can be further increased by hydrolyzing the fiber portions of CGM or by increasing the glucose content by addition of other polysaccharide depolymerase enzymes. The pH, temperature and time are chosen to achieve the optimal enzyme activity using known techniques and specifications.

The destarched corn gluten is then treated with a mild alkaline buffer (i.e., pH 7.5 to 9) to improve removal of fatty acids and residual corn oils from the enzyme hydrolyzed corn gluten. Alkaline treatment further aids in removal of compounds from the corn gluten

-7-

which impart its characteristic strong, bitter off-flavor. This step is performed using sodium or potassium carbonate/bicarbonate at a pH of from about 8 to about 9 and a temperature of from about 60 to about 80°C, preferably 80°C. Other mild alkaline solutions can be used such as sodium hydroxide, calcium hydroxide or potassium hydroxide, provided that the pH does not exceed about 9. Very little or substantially none of the xanthophyll, zeaxanthophyll and carotene natural food pigments are solubilized at this stage. This permits subsequent alcohol extraction and recovery of these pigments under conditions which insure their native structure, minimize high alkaline-induced by-product formation (i.e., pH>9) and minimize zein solubilization. Aqueous washing between the enzymatic hydrolysis and alkaline treatment steps removes most of the carbohydrate that may form flavor and color compounds. After alkaline treatment, the pH can be lowered (e.g., from about pH 6.5 to about 7.5) by adding a suitable organic or inorganic acid (e.g., acetic acid or HCl) with good mixing.

After enzyme hydrolysis and alkaline treatment, the resulting gluten will still contain glutelin(s), pigments, zein(s) and various hydrocarbons. This form of gluten is then decolorized and further deflavored by a series of alcohol washes to harvest natural food colorants and simultaneously remove minor residual off-flavor from the gluten. The concentration of alcohol is chosen to specifically solubilize pigments and flavoring agents but not substantially solubilize the zein portion. The solubility of a given zein component of CGM can be ascertained by selecting conditions (i.e., alcohol concentration and temperature) at which zein is not

-8-

extracted along with the impurities. See Figure 2 which graphically illustrates how alcohol temperature and titer affect zein solubility. As shown in this Figure, room temperature (about 25°C neutral pH i.e., pH 6.5 - pH 7.5) alcohol having a titer of from about 10 to about 40% (low titer alcohol) or from about 90 to about 100% alcohol (high titer alcohol), can be used for washing CGM without extracting zein. In each of these ranges, zein is substantially insoluble. One of the advantages of this invention is that brief exposure to intermediate titers (i.e., 40-90%) can occur without substantial loss of zein. High titer alcohol washes are preferred since the quantity of alcohol required to remove colors and impurities is minimized, due to the increased solubility of the impurities. Low titers of alcohol can be used but larger quantities of alcohol are required to adequately extract the impurities. However, one advantage of using low titer alcohol is that due to the large volume of water for dilution, it is possible to reduce and/or to eliminate subsequent water washes.

The conditions which affect zein solubility in various solvents have also been described in detail by Dill, J. Biol. Chem. 77:239-247 (1927) and Manley, Indus. Eng. Chem. 35:661-665 (1943). The teachings of these papers are incorporated herein by reference.

During the pigment extraction step, zein(s) remains within the zein body membrane, which protects the alpha zein (soluble in high titer alcohol(s) such as isopropanol/ethanol) and beta zein (soluble in lower titer alcohol) fractions from being extracted along with the pigments. Under optimal conditions, zein remains encapsulated in the zein body membrane, thereby preventing zein loss. Thus, the xanthophyll/alcohol extract

-9-

will remain essentially uncontaminated with zein(s) during the alcoholic extractions.

It is desirable to use high titer alcohol (i.e., 90 to 100%) for the depigmentation/deflavoring step to avoid solubilizing the zein during the extraction of pigments and contaminants. It should be noted that as the concentration of alcohol decreases (i.e., falls inside each of the curves shown in Figure 2), the solubility of zein increases, thus resulting in the extraction of zein and overall loss of product. Preferably, alcohol washing is conducted at ambient temperature or at slightly elevated temperature. The preferred method of washing is in a continuous counter-current fashion or by batch-wise washing with fresh alcohol at each step. Either ethanol or isopropanol are the preferred solvents but can be replaced with other alcohols and food grade solvents that are miscible with water and have a limited titer range for which zein is soluble. A solvent and concentration range can be selected at which pigments and contaminants are maximally soluble and the zein is negligible or minimally soluble.

By-products of the depigmentation step, particularly xanthophyll - a yellow pigment, can be further purified using standard techniques to yield food grade food colors. Purification of the pigments can be accomplished using ion exchange resins, for example.

The xanthophyll and carotene-free corn gluten is then passed through a series of water washes to remove water-soluble products trapped by residual pigments/hydrocarbons which may not have been removed by previous wash steps, enzymatic hydrolysis and alkaline treatment of the gluten. Residual alcohol is also removed from the gluten by the water washes. The number of water washes

-10-

depends upon the degree of purity desired. This can be assessed by ascertaining the amount of residual water-extractable flavor compounds removed after each washing. It should be noted that between the alcohol and water wash steps, the destarched and decolored gluten is recovered by centrifugation, filtration or freeze drying.

An alcohol extraction step is then performed to separate the gluten into a zein fraction and a protein fraction containing glutelin. During this extraction step, it is necessary to choose a concentration of alcohol and temperature at which zein is readily soluble. The preferred solvents for zein are aqueous solutions of ethanol, 2-propanol or mixtures thereof, which contain at least 40%, but less than, 90% alcohol, or the equivalent range for isopropanol. Commercial zein is typically soluble in 80:20 mixtures of 2-propanol/water in amounts up to at least 30% (v/v) protein, over a temperature range of about 5-70°C; and in 80:20 ethanol/water in up to at least 30% (v/v) protein, over a temperature range of about 5-70°C, taking into account that the temperature should remain at least 5°C below the boiling point of the solvent. For some zein applications, like film formation, it is preferable to select an alcohol titer at the upper end of the 40-90% range to minimize isolation of beta zein and glutelins and to maximize alpha zein extraction and purity. For some other applications where maximizing zein yield is most important, extractions at both high and low titers can be made and later recombined as necessary.

The yield of zein extracted can be optimized at elevated temperature (i.e., from about 25-70°C, with 50°C preferred) in order to facilitate extraction of the zein

-11-

from remaining glutelins and zein body membrane degradation products. It is also necessary to adjust the pH of the alcoholic solution to below pH 7.0 and preferably within the range of from about 5.5 to about 6.5 to improve zein solubility. Once the pH of the ethanolic solution has been adjusted, the pH should be maintained near the isoelectric point for most zein phenotypes (i.e., from about pH 6 to about 7) to minimize extraction of marginally soluble zein(s).

In a preferred embodiment, a solution of aqueous ethanol is used containing from about 70 to about 90% ethanol; that is, having a water:ethanol ratio of from about 30:70 to about 10:90. Preferably, aqueous 80% ethanol (vol) can be used for the extraction with high yield of zein.

The glutelin by-product can be recovered from the extraction cake and further purified using known techniques for use as a vegetable protein supplement. In order to achieve maximum solubility of the resultant glutelin fraction, the last zein extraction should be followed immediately by equilibrating the glutelin in <50% ethanol containing >1% humectant (based on the glutelin dry weight) chosen from the following non-limiting list or their analogs (glycerol, propylene glycol, polyethylene glycol (PEG), etc.), <2.5% gum chosen from the following list (xanthan, gelatin, konjac, alginate or its derivatives, cellulose or its derivatives, carrageenans, starch or its derivatives, locust bean gum or guar gum). Especially preferred, are the gums, like xanthan, hydroxypropylated alginate and cellulose which are soluble in alcohol at titers between 30 and 50%, i.e., which coagulate other gums and thereby inhibit uniform penetration and mixing with the glutelins. It is

-12-

also desirable to incorporate a small amount (<1.25%) of food grade emulsifier into the glutelin fraction. The selection of a specific blend of these protein stabilizers will depend on the intended food application of the glutelin as a vegetable protein supplement. After formulation with the above protein stabilizers, it is preferable to dry the glutelin by-product by freeze drying. Other means for drying the product include spray drying and atmospheric drying at <50°C.

Zein can then be recovered from the alcoholic solvent and concentrated using standard techniques, such as but not limited to precipitation, ultrafiltration, evaporation or other appropriate techniques, or reduced to a dry powder by flash drying, lyophilization or spray drying. The preferred method for obtaining a readily dispersible powder is either to reduce the ethanol titer to about 15% and spray dry, or to disperse the ethanolic zein solution in a slightly alkaline (pH about 7 to 8.5), well-stirred, aqueous medium followed by gentle (<50°C drum or standard freeze drying) dehydration to yield zein containing <10% water.

In one embodiment, zein can be further purified, depending upon end use, prior to recovering it from the alcoholic solution by passing the protein solution through a column, or an adsorbent, such as activated carbon (granular or powdered forms) and macroreticulated resins, such as ion exchange and adsorption resins.

The purified zein produced by the methods of this invention is essentially decolorized (i.e., the product is white), odor-free and deflavored. The decolorized and deflavored zein is also substantially free of hydrocarbon extracts which are otherwise present using other purification techniques. Thus, purified zein produced according

-13-

to the methods of this invention is suitable for use in foods and pharmaceutical applications, such as coatings, texturizers, additives and fat replacers. For example, an aqueous dispersion of deflavored and decolored zein can be cast into a continuous edible film having suitable strength and protective barrier properties. Alternatively, a film can be cast from an ethanolic solution to produce an edible barrier coating.

Prior to recovery from the alcoholic solvent, the decolored and deflavored zein can be further processed. In a preferred method, the ethanolic zein solution is injected into stirred water containing an anti-aggregate to produce water-dispersible microparticles of hydrophobic, water-insoluble zein. Such techniques are described in U.S. Patent 5,021,248 (Stark and Gross), the teachings of which are incorporated herein by reference. In another embodiment, the ethanolic zein solution is dispersed as a mist before it is contacted with water, or as part of co-focused mist containing ethanolic zein and aqueous antiaggregate solutions. In each of these processes, gums and/or additive can be added to the ethanolic zein solution depending upon the desired end use.

In another embodiment of the invention, deflavored and decolored gluten, obtained after depigmentation with alcohol, can be harvested for use in foods, without additional purification or separation. For example, a concentration of 4% gluten or zein bodies (w/v) can substantially increase (>30%) the baked volume of a microwave cake without compromising its texture or taste. Deflavored and decolorized gluten also exhibits fat mimetic properties of microparticulated proteins, such as those described in U.S. Patent No. 5,021,248, and can be

-14-

used as a fat replacer and texturizer. The decolorized and deflavored gluten can be produced which contains an enriched zein body fraction (i.e., >75%). The optimum proportion of zein bodies to glutelin microparticles is dependent on the intended application. In fat replacer applications like frozen desserts, mayonnaise and whipped topping, for example, it is desirable to have an enriched gluten having >90% zein bodies. For fat replacement in sausages, however, a 50-75% incidence of the zein body fraction tends to be adequate. For microwave cake volume expansion, only a 50:50 mixture of zein bodies to glutelins produces the 30% volume expansion functionality without compromising taste or texture. Other proteins, including wheat gluten, do not have this effect. The fractionation of zein bodies and glutelin microparticles can be achieved by low speed centrifugation in alcohol, which contains aqueous alcohol soluble gum(s) and/or aqueous alcohol soluble salt(s) to effect the desired degree of zein-body enrichment. In addition, it is possible to program the sieving and gelation of the centrifugation medium by the use of a thermally reversible gelling agent (<4%) to effect the removal of the largest particles (i.e., those most likely to cause a gritty or chalky mouthfeel). The simplest application of the latter technique is to centrifuge at a temperature which maintains a relatively low viscosity (i.e., 1-400 m.Pa.S) during the centrifugal separation of the two fractions followed by lowering the temperature to below the gelling point to effect gelation and thereby prevent component mixing in the absence of a density gradient. After separation and gelation, the centrifugation medium can then be sectioned and the zein body-rich fraction and glutelin-

-15-

rich fraction recovered by reliquefaction (<50°C) of the dilute gel medium. This technique is facilitated, in part, by the fact that zein is a very robust protein with respect to heat and is not readily denatured even by high temperatures (i.e., 125°C) in glycerol.

The invention is further illustrated by the following non-limiting examples.

EXAMPLES

EXAMPLE 1 PURIFICATION OF ZEIN

A slurry was prepared by mixing 3.3 L of deionized water with 2.8 kg of wet corn gluten meal (CGM) from representative corn wet-milling plants (Grain Processors Corp.; Muscatine, IA or Corn Products Corp., Argo, IL). The gluten is nominally 40% solids and is collected before it was dried or admixed with other process streams. The slurry was vigorously agitated with a Lightnin[®] Mixer (Model TS1515, General Signal Corp., Mixing Equipment Div., USA) using a 3-inch diameter impeller to uniformly disperse the gluten. After heating the slurry to 80°C, the pH was adjusted to 6.4±0.1 using 3N sodium hydroxide.

To digest the starch, 5.1 mL of α -amylase was added to the wet gluten slurry (Taka-Therm[®] L-340, Solvay Enzyme Products, Inc., Elkhart, IN), reacted for 1 hour and then cooled to 50°C. The slurry was centrifuged in a solid bowl centrifuge at 1200xg for 10 min and the supernatant discarded. The pellet was resuspended in an equal part of hot water (80°C) and centrifuged again at 1200xg. The supernatant was discarded. No substantial reduction in off-flavor was detected when the destarched gluten was tasted.

-16-

The destarched gluten pellet was then treated with 1M sodium carbonate and 1M sodium hydrogen carbonate to adjust pH to 9.0 at 20°C. The slurry was then heated to 80°C and found to initially produce an acrid, unpleasant odor. However, after 1 hour of agitation the headspace odor changed to a pleasant, almost sweet-smelling cereal odor. At the end of that time, the slurry was cooled to 50°C. The slurry was centrifuged in a solid bowl centrifuge at 1200xg for 20 minutes and the supernatant discarded. The pellet was resuspended in an equal part of water and neutralized with 3N HCl. The solids were centrifuged two additional times and resuspended in an equal part of water. Even though the gluten was still bright yellow, there was no longer any offensive bitter off-taste.

The centrifuged pellet containing 599.5 g of solids was reslurried in 3 volumes of 100% (neat) ethanol at room temperature for 15 minutes. The slurry was centrifuged and the alcoholic supernatant discarded. The centrifuged pellet was resuspended five times in 2 volumes of 100% (neat) ethanol for 40 minutes. The first and second supernatants show minimal zein losses. Zein was not detected in the liquor of the third through sixth supernatants. The pellet after the sixth ethanol wash yielded 512.3 g of solids, for a loss of 11.6% throughout the ethanol washing. The successively higher alcohol titer after each wash is summarized below in Table 1.

-17-

TABLE 1

<u>Ethanol Wash</u>	<u>Estimated Ethanol Titers</u>	<u>%Zein* Loss</u>
1	77.3% (wt)	2.3
2	93.1%	0.4
3	97.9%	-
4	99.4%	-
5	99.8%	-
6	99.9%	-

*Based on an estimated 32% zein content in the initial CGM, zein loss was determined by dropping some ethanolic wash solution into a large volume of water. Any cloud formation in the water indicated the presence of zein. It was further quantitated by measuring HPLC peak areas.

The pellet was then washed three times with 2 volumes of water and centrifuged. A small amount of zein was detected in the first and second wash waters. The total zein loss was 1.6% of the zein in the CGM starting material. The CGM cake, at this point, was grey, having lost all traces of yellow color. The fully decolorized gluten was found to have no objectionable off-taste. However, a very slight residual cereal taste was still detectable and the need for its complete removal for some applications, was recognized. All washes and centrifugations were carried out at room temperature.

The water washed pellet was resuspended in an alcoholic solution calculated to raise the ethanol level in the final slurry to 80% (v/v) ethanol and permit complete dispersion of the solids. This number is a function of the pellet dry solids content. The amount of neat ethanol added to the water washed pellet was determined using the following equation:

-18-

Let X = volume of 100% ethanol, mL

W = wt % of pellet

Y = weight of pellet, g

$$X = 3.476 * (1 - (W/100)) * Y$$

If 95% ethanol was used, the equation was modified to:

Let X = volume of 95% ethanol, mL

$$X = 4.661 * (1 - (W/100)) * Y.$$

The slurry was stirred and heated to 55-60°C, the pH was adjusted to 5.9±0.1 using 1 N HCl, and the extraction of zein proceeded for one hour. The resulting slurry was centrifuged and the supernatant containing zein was collected. At this stage, the pellet can be subjected to additional extractions to recover residual zein. However, it has been observed that subsequent extractions of the pellet contained a higher proportion of off-flavor compounds and required further processing to remove them.

Zein can then be recovered from the alcoholic supernatant by adding enough water to lower the final ethanol titer to less than 40% and cooling to room temperature. The resultant zein precipitate can then be recovered, dried, and ground to yield a slightly, off white, almost fully deflavored corn protein powder.

EXAMPLE 2 Further Purification Using Activated Carbon

Example 1 was repeated but before the zein was precipitated from the alcoholic suspension, it was further purified using activated carbon. Powdered carbon (Darco SGL80x325, American Norit Co., Inc., Jacksonville, FL) was added to the ethanolic zein liquor at 15% of the

-19-

zein content. This treatment was performed at ambient temperature for one hour. The carbon powder was then filtered out using a scintered glass filter funnel and a bed of filter aid (Celite® 521, Aldrich Chemical Co., Inc., Milwaukee, WI). The resulting zein powder was whiter and more deflavored, after precipitation and drying than the product produced by the methods of Example 1.

EXAMPLE 3 DESTARCHING

Method 1

An α -amylase to gluten ratio different than that of Example 1 was used to effect the hydrolysis of starch in CGM. A slurry was prepared by mixing 2.4 kg of CGM with 3.3 L of deionized water. The slurry was vigorously agitated to uniformly disperse the gluten. After heating the slurry to 80°C, the pH was adjusted to 6.4 \pm 0.1 using 3N sodium hydroxide.

To digest the starch, 5.1 mL of α -amylase solution was added to the slurry (Taka-Therm® L-340, Solvay Enzyme Products, Inc., Elkhart, IN). Reaction time was 1 hour, after which the slurry was cooled to 50°C. Progress of the reaction was followed by periodically removing aliquots of the slurry and using an iodine indicator. Starch will form a blue or purple complex with the iodine according to the classical procedure (AOAC 925.56). At the end of one hour, there was no trace of iodine-starch complex present in the slurry.

Method 2

After starch hydrolysis according to Example 1, the slurry was centrifuged twice at 1200 xg for 20 minutes

-20-

and reslurried twice with an equal part of water. All steps were carried out at ambient temperature. No substantial reduction in off-flavor was detected.

Method 3

After starch hydrolysis according to Example 1, the slurry was centrifuged at 1200 xg for 20 minutes and reslurried three times with an equal part of water. All steps were carried out at ambient temperature. No substantial reduction in off-flavor was detected.

Method 4

A slurry was prepared by mixing 1.1 kg of CGM with 2.2 L of deionized water. The starch was removed from the gluten according to the method described in Example 1. The water wash after starch digestion was not performed. Instead, the slurry was centrifuged and prepared for carbonate treatment.

Destarched gluten slurry amounting to 2.4 kg was further diluted with 2 L of deionized water. To this slurry, 240 mL of 1 M sodium hydrogen carbonate and 130 mL 1 M sodium carbonate were added and adjusted to pH 9.3. The slurry was then heated to 80°C. It was held at that temperature under agitation for 1 hour. At the end of that time, the slurry was cooled to 50°C. The slurry was centrifuged at 1200 xg for 20 minutes and the supernatant discarded. The pellet was resuspended in an equal part of water and neutralized with 3N HCl. The solids were twice more centrifuged and resuspended in an equal part of water. All centrifugations were performed at ambient temperature.

Centrifuged pellet containing 654.8 g of solids was reslurried in 2 volumes of 100% (neat) ethanol at ambient

-21-

temperature for 15 minutes. The slurry was centrifuged and the alcoholic supernatant discarded. The pellet was resuspended seven times in 2 volumes of 100% (neat) ethanol. Each time the suspension was centrifuged, the supernatant was discarded. Two more ethanol washes were required to get to the same gray pellet color as observed using the methods of Example 1.

EXAMPLE 4 CARBONATE TREATMENT OF DESTARCHED WET GLUTEN

Method 1

Wet corn gluten was destarched according to Example 1. The destarched slurry was treated with 1M sodium carbonate and 1M sodium hydrogen carbonate to adjust pH to 8.7 ± 0.1 . The slurry was heated to 80°C for 1 hr while agitating. The slurry was centrifuged and the supernatant discarded. The pellet was twice suspended with an equal part of water and centrifuged.

Method 2

The effect of ethanol versus water washes on the removal of off-color was evaluated. Destarched gluten was carbonate treated with 1000 mL of 10% 0.1 M sodium carbonate and 90% 0.1 M sodium hydrogen carbonate at 9 pH. The mixture was stirred for 1 hr at 80°C and then cooled to room temperature. The mixture was maintained at room temperature and agitated for 3 days. The mixture was split into equal portions. The first portion was washed three times with 50% ethanol and 0.2% Tween 80. The second portion was washed three times with water. The water washed gluten, which was not exposed to either alcohol or surfactant, was much lighter in color than the (i.e., ethanol/Tween 80) ethanol/Tween 80 washed gluten

-22-

(which was dark greenish brown) and also much less yellow than the destarched but not carbonated treated gluten. Water washing was therefore considered to be highly preferable to more expensive and complex methods involving alcohol and surfactants.

EXAMPLE 5 ETHANOL WASHING OF DESTARCHED, CARBONATE
TREATED GLUTEN

Method 1

1100 mL of carbonate-treated, water-washed CGM (15% solids) prepared by the methods described in Example 1 was centrifuged, lyophilized, and resuspended in 1320 mL of absolute ethanol. The suspension was homogenized using a Polytron homogenizer to disperse the solids. The dispersion was centrifuged and the pellet was resuspended in 100% (neat) ethanol to the amount of 2 volumes of original dried cake (165 g). The alcohol washing step was repeated 6 times with 200 mL of absolute ethanol for each wash. All washes were conducted at room temperature. The ethanol washes were followed by two water washes and then the gray colored cake was reslurried in water and centrifuged at 3,000xg for 8 min. The resultant cake was still gray but represented decolorized and deflavored gluten which is suitable for some food applications.

Method 2

1 Kg of CGM was carbonate-treated, water-washed, centrifuged, and the pellet was reslurried in 3 volumes of 95% ethanol at room temperature for 2 hours. Zein loss was determined in the manner described in Example 1. Zein was not detected in any of the supernatant ethanolic solutions, although substantial yellow pigment removal was evident. The alcoholic supernatants were

-23-

then set aside for later xanthophyll pigment recovery. The centrifuge pellet was resuspended in 3 volumes of 70% ethanol for 30 minutes. Again, zein was not detected in the supernatant liquor. After centrifuging, the pellet was resuspended three more times in 1 volume of 70% ethanol. Finally, the centrifuge pellet was washed three times with 1 volume of water and centrifuged. The estimated ethanol titers during the first five washes are listed below:

<u>Ethanol Wash</u>	<u>Estimated Ethanol Titters</u>
1	78.6% (vol)
2	71.6%
3	70%
4	70%
5	70%

Method 3

Destarched, carbonate-treated CGM produced by the methods described in Example 1, was divided into two portions. The first portion (25g of a 29.6% solids slurry) was washed twice with 3 parts 100% (neat) ethanol. Fresh alcohol was used each time. The supernatant from the second wash was used to wash the second portion of CGM (19.3g of the slurry). In this way, a two stage counter-current extraction was evaluated. Ethanol liquors from all three washes were analyzed by gel permeation high performance liquid chromatography (HPLC) according to the methods described below.

-24-

Solvent System: Isocratic
60% H₂O/40% acetonitrile/0.1% phosphoric
acid
40 minute run

Column: Bio-Sil TSK-250 600 x 7.5 mm (Bio Rad)
Bio-Rad Gel Filtration Column (Catalog#125-0066)

Program: End time - 38 min
Narrowest Peak - 1.4-3.2 sec
Peak Width - Automatic
Base Treatment - B to B
Area Sensitivity - 1
Base Sensitivity - 1
Skim Sensitivity - 1
Area Rejection - either 1×10^5 or 1×10^4

Detection: UV λ = 214nm

Dilutions: For ethanolic zein extracts and ethanol
gluten washes: A 1:1 with solvent
For UF washes - no dilution
AZTL Procedure - no dilution

Injection volume: 100ul

The peak areas for the counter-current sample showed about a 96% zein recovery efficiency compared to the combined peak areas from the first and second fresh alcohol washes. A Summary of the alcohol washes is shown in Table 2.

-25-

TABLE 2
Integration Area X 10⁻⁶

	<u>Fresh</u> <u>EtOH</u>	<u>Fresh</u> <u>EtOH</u>	<u>Fresh</u> <u>EtOH</u> 1st & 2nd Wash	<u>Counter-</u> <u>Current</u> <u>Wash</u>
	<u>1st Wash</u> <u>Column#1</u>	<u>2nd Wash</u> <u>Column#2</u>		
Protein Area	43.617	.011	43.628	55.566
Total Non-				
Protein Area	66.052	12.809	78.861	76.004
Total Area	109.669	12.820	122.489	131.570

Counter Current =	Total-Protein Area For Counter Current Wash	=	76.00	=	96.3%
Wash Efficiency	Total Protein Area for Fresh Wash		78.86		

EXAMPLE 6 EFFECT OF TEMPERATURE ON COLOR REMOVAL

50 ml of destarched and carbonate-treated gluten cake prepared by the methods described in Example 1 (8% solids), was divided into two portions. One portion was washed (5 min) five times (40 ml each) in 100% ethanol at 25°C. The other portion was washed similarly at elevated temperature (35-55°C). The slurries were centrifuged at 2000 rpm for 10 minutes. The supernatants were sampled and absorbance at 450 nm measured to determine the amount of pigments extracted from the gluten cake. A final wash was done with water after the level of pigments detected in the ethanolic wash supernatant became insignificant.

The results of this pigment exaction are shown in Figure 3 and indicate that the elevated ethanol temperature did not significantly improve the amount of pigment

-26-

extracted from the gluten. The quantity of extracted pigment was essentially the same using 23°C ethanol and 35-55°C ethanol.

EXAMPLE 7 PREPARATION OF FILM FORMING BARRIER

A zein film (9 x 14" x 4mil) was cast by solvent evaporation (50°C) from 150 mL of a 6% commercial zein (Freeman Industries, Inc., yellow granular powder) solution (80% ethanol) because a 4% solution did not adequately produce a continuous film. In contrast, only a 4% zein film cast under the same conditions from white zein made by the methods of this invention was continuous, strong and could be bent without breaking. The O₂, CO₂ and H₂O barrier properties of the two films were evaluated using a Mocon analyzer (Modern Controls Inc., Minneapolis, MN). The results of this study are shown in Table 3.

-27-

TABLE 3

Gas	Film Thickness (mil)	Transmission Rate		OPTAZein Advantage ⁴
		cc/100 in ² /24 hrs	Permeance ³ (Ave.)	
H ₂ O OZ ¹	2.0	10.1	20.5	32%
	2.1	9.9		
	FZ ²	5.1	30.2	
	3.9	8.3		
CO ₂ OZ	4.1	3.1	22.3	53%
	2.0	16.0		
	FZ	6.3	47.7	
	3.4	18.8		
O ₂ OZ	3.9	5.5	21.9	26%
	2.1	10.7		
	FZ	4.0	29.6	
	3.2	11.9		

1. OZ = OPTAZein produced by methods of this invention
2. FZ = Commercial Zein 4000, Freeman Industries, Inc.
3. Permeance determined by methods of (ASTM F-1249-90 and ASTM D-3985-81) at 23°C.
4. The better barrier properties of the zein produced by the methods of this invention are expressed as a % reduction in permeance relative to commercial Freeman zein 4000.

$$\left(\text{i.e., } \frac{\text{FZ-OZ}}{\text{FZ}} \times 100 = \% \text{ OPTAZein Advantage} \right)$$

The results above indicate that the film cast from 4% zein produced by the methods of this invention served

-28-

as a better barrier than films cast from 6% commercial zein. It should be noted that the film produced using commercially available zein was yellow in color, in contrast to the OPTA zein film which was transparent.

EXAMPLE 8 COMPARATIVE ANALYSIS

The taste of the zein produced, according to the methods of this invention, was assessed by both high performance liquid chromatography (HPLC, technique described earlier) and a sensory panel using the aqueous fractions of 80% ethanolic solutions made from each type of zein. The aqueous extract was obtained by selectively evaporating the ethanol under vacuum to concentrate the water soluble organic constituents. The resultant aqueous fraction chromatograms are shown in Figure 4.

The proximate analyses of wet gluten, destarched wet gluten, deflavored, decolored zein (hereinafter "dfdc zein") and deflavored, decolored glutelin (hereinafter "dfdc glutelin") (on a dry solids basis) and particle size distribution throughout this process are shown in Table 4.

TABLE 4
Proximate Analysis¹

	Wet	Destarched	dfdc ²	dfdc Glutelin
	Gluten ²	Wet Gluten ²	Zein	Cake After Zein Extraction ²
Protein	70.2%	92.1%	96.98	93.0%
Fat	4.8%	1.6%	<0.1	0.5%
Ash	1.3%	0.5%	0.5	0.4%
Total				
Carbohydrate	23.7%	5.8%	2.51	6.0%
Dietary Fiber	8.1%	5.8%	n.d. ³	n.d. ³

1. A.O.A.C., 15th Edition Method 981.10 (1990).
2. determined on an anhydrous basis
3. n.d. = not determined

Table 4 illustrates the increasing percentage of protein typically achieved by the invention as one proceeds from the wet gluten starting material to the intermediate destarched wet gluten and finally to the dfdc zein and dfdc glutelin. It should be noted that the standard proximate analysis technique does not specifically determine carbohydrate (i.e., using phenol, anthrone or other carbohydrate-specific reagents. Instead, the protein nitrogen was specifically determined using a micro Kjeldahl technique. Kjeldahl Test, pg 674, Hawley's Condensed Chem. Dictionary, 11th Ed. (1987) Van Nostrams, Reinhold, New York. The fat was determined by extraction in ether and the ash content was determined gravimetrically using an ashing furnace. The appropriate factor for nitrogen in the micro Kjeldahl calculation to convert to % protein is a function of the amino acid

-30-

content of the protein sample. When analyzing for the largest component in the sample (i.e., protein), small changes in the % nitrogen to % protein conversion factor can make significant differences in the protein purity determination. For example, in the case of dfdc zein, the ash (which was specifically determined), and the fat (which was determined by extraction) are $\leq 0.6\%$ by weight of the sample, it is possible that the protein content could be 99.4% (i.e., adding 2.51% to 96.98%) instead of 96.08% depending upon the conversion factor used. Such a conclusion would require a precise determination of the sample's amino acid content and appropriate change in the Kjeldahl nitrogen factor.

EXAMPLE 9 ZEIN BODY AND GLUTELIN MICROPARTICLE ISOLATION

One of the principal advantages of the invention is that it preserves the natural microparticulation of its zein body and glutelin components as shown in the "Median Size" Column in Table 5 below:

-31-

TABLE 5

<u>Process Step</u>	<u>Mean Size (μm)</u>	<u>Median Size (μm)</u>	<u>10% Size (μm)</u>	<u>90% Size (μm)</u>
Feed Gluten	15.86	10.29	4.87	35.25
After Amylase	13.85	8.48	3.93	31.32
After Carbonate	8.97	6.17	2.95	15.41
After 1st EtOH Wash	17.45	8.43	3.85	45.68
After 2nd EtOH Wash	13.1	7.63	3.40	31.09
After 6th EtOH Wash	21.21	8.75	3.47	60.98
After 3rd H ₂ O Wash	23.30	9.92	3.76	66.30

If the median particle size is used as an index, then the original gluten particle size distribution is essentially unchanged during processing. The rise in mean size and 90 percentile values is due to some agglomeration.

After isolation of the dfdc gluten, in either wet or gently dried forms (preferably containing a humectant), the zein body and glutelin components can be isolated by centrifugation using one of the methods below. The specific method chosen will typically depend more on the intended food application than on any other factor.

The isolation of zein bodies from native corn endosperm - as opposed to wet milled gluten, has been reported using 15% sucrose gradients which are impractical for commercial use. Instead, it has now been discovered that the centrifugal isolation of zein body-rich and glutelin-rich fractions which meet commercial food needs

-32-

is possible using a combination of viscosifying gums and salting out techniques. Method 1 illustrates the principle for aqueous separations while Method 2 illustrates a conceptually similar separation in an alcohol solvent of varying titer.

Method 1 Aqueous Dispersion of dfdc Gluten

A 20% (w/v) suspension (25°C) of dfdc gluten was prepared in water or taken as an appropriate dilution of the last water wash in Example 1. It was mixed 1:1 with a 2% suspension of a suitable gum (listed earlier) and cooled after dissolving to 25°C. Thus, 10g (dry solids equivalent) of dfdc gluten was suspended in 50ml of distilled or deionized water and homogenized briefly (10,000rpm, 1 min) in a Polytron homogenator. The resultant suspension was combined with 50ml of 2% xanthan gum in water and the two are mixed at low speed (<500 rpm) for 5 min. The gluten suspension was divided between two 50mL plastic centrifuge tubes and centrifuged for 5 min at approx. 2000 rpm and the relative enrichment of the two fractions was monitored using a microscope (1200 x magnification). This non-equilibrium separation permitted recovery of a zein body-rich fraction ($\geq 75\%$ zein bodies in the supernatant and a glutelin-rich fraction in the precipitated or heavier fraction). If the separation is inadequate, additional centrifugation under the above conditions can be done. Addition of divalent salts (e.g., calcium salts), adjustment of the pH to the pI of the glutelin microparticles (determined by zein potential measurement in free-flowing electrophoresis), and/or ammonium sulfate precipitation can further facilitate isolation or fraction-enrichment.

Method 2 Ethanolic Dispersion of dfdc Gluten

Since there are alcohol titers (and pHs) which zein is not extracted from the zein bodies, it is possible to separate the two fractions by centrifugation in various titers of alcohol. When 70-100% titers of alcohol are used, it is important to add some suitable humectant (<1%; described above) to prevent dehydration of the protein, aggregation and other adverse textural effects.

A 10% (w/v) suspension of dfdc gluten was prepared in either high titer ethanol (85-100%) or low titer ethanol (0-55%). In this instance, a 95% ethanol titer was used containing 2% CaOAC (which is an alcohol soluble salt) and 5% glycerol, 50 ml of the CaOAC/glycerol/xanthan-containing ethanol was added to the 50mL of 20% gluten suspension and centrifuged for 10 min. at 50 rpm. The resultant separation was monitored microscopically (800-1000X) and additional centrifugations were performed if the incidence of zein body purity was less than desired.

The fractions were separated by decanting the light zein body-rich supernatant which represents approximately 50% of the dfdc gluten weight. Since the zein body fraction, thus isolated appears to be 95% pure, this represents a 50% yield of the total zein bodies in the gluten. Since such separations are possible in both aqueous and alcoholic fluids, it is likely that an analogous separation could be effected in a suitable air classification apparatus (i.e., hydroclone, etc.).

-34-

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims:

-35-

CLAIMS

1. A method of producing decolored and deflavored zein or zein bodies and glutelins from corn gluten comprising removing color and flavor impurities from the gluten by treating corn gluten with alkali to remove fatty acids and corn oils contained therein and then with an aqueous alcoholic solvent having a concentration which does not substantially extract zein, zein bodies and glutelins therefrom, thereafter changing the concentration of the alcoholic solvent to a concentration sufficient to separate the gluten into decolored and deflavored zein or zein bodies and glutelin fractions.
2. The method of Claim 1, wherein the alcohol is a food grade alcohol.
3. The method of Claim 2, wherein the alcohol is ethanol, isopropanol or combinations thereof.
4. The method of Claim 1, wherein the alcohol concentration is of from about 90 to about 100% by weight or from about 10 to 40% by weight alcohol and is then changed to a concentration of from about 40 to about 90% by weight alcohol.
5. An edible, essentially decolored and deflavored zein produced by the method of Claim 1.
6. Essentially decolored and deflavored zein bodies that are edible produced by the method of Claim 1.

-36-

7. A method of producing decolored and deflavored zein or zein bodies containing zein therein from corn gluten, comprising the steps of:
- a) destarching the corn gluten with a carbohydrate hydrolyzing enzyme under agitating conditions, sufficient to hydrolyze and remove starch contained therein;
 - b) treating the destarched corn gluten of step (a) with an alkaline buffer having a pH from about 7.5 to about 9 under conditions sufficient to remove fatty acids or corn oils contained therein;
 - c) selectively extracting pigments and flavor compounds from the destarched and alkaline treated gluten of step (b) with aqueous alcohol having an alcohol concentration that does not substantially extract zein, zein bodies or glutelins;
 - d) separating the gluten into zein or zein body and glutelin fractions by extracting or fractionating the gluten of step (c) with aqueous alcohol solvent having an alcohol concentration sufficient to either solubilize the zein or fractionate zein bodies therein; and
 - e) recovering the separated zein or zein body fractions from the appropriate aqueous alcoholic medium to thereby yield decolored and deflavored zein or zein bodies containing zein therein.
8. The method of Claim 7, wherein the enzyme is selected from the group consisting of α -amylase, pullulanase, cellulase and glucoamylase.

-37-

9. The method of Claim 7, wherein step (b) is performed at a pH of from about 8 to about 9 and a temperature of from about 60 to about 80°C.
10. The method of Claim 7, wherein the alkaline buffer is selected from aqueous solutions of sodium hydroxide, calcium hydroxide, ammonium hydroxide, sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate and mixtures thereof.
11. The method of Claim 7, wherein step (c) is performed with ethanol or isopropanol having an alcohol concentration of from about 90 to about 100% by weight or from about 10 to about 40% by weight.
12. The method of Claim 7, wherein step (d) is performed using an ethanol concentration of from either about 40 to from about 90% by weight alcohol or from about 0-40% by weight alcohol.
13. The method of Claim 7, wherein step (e) is performed by precipitating the purified zein in water or from about 10 to about 40% by weight alcohol, removing the zein by filtration therefrom and then drying the product.
14. The method of Claim 7, further comprising washing the product of step (b) with water to remove residual water soluble products from the gluten.
15. The method of Claim 7, further comprising the step of treating the extracted zein with activated carbon or ion exchange resin to remove impurities in the alcohol extracted zein.

-38-

16. An edible, essentially decolored and deflavored zein purified by the method of Claim 7.
17. An edible, continuous film cast from an aqueous solution containing decolored and deflavored zein purified by the method of Claim 7.
18. An edible, essentially decolored and deflavored zein body fraction purified by the method of Claim 7.
19. The method of Claim 7, further comprising recovering the glutelin fraction to thereby yield decolored and deflavored glutelins.
20. An ethanol solution of edible, deflavored and decolored zein, purified by the method of Claim 7, which is suitable for use as an edible barrier coating.
21. An essentially decolored and deflavored zein that is suitable for use in a food or pharmaceutical application.
22. A food product containing decolored and deflavored zein, zein bodies, glutelins and combinations thereof.
23. A method of producing deflavored and decolored corn glutens, comprising the steps of:
 - a) destarching the corn gluten with a carbohydrate hydrolyzing enzyme under agitating conditions, sufficient to hydrolyze and remove starch contained therein;

-39-

- b) treating the destarched corn gluten of step (a) with an alkaline buffer having a pH from about 7.5 to about 9 under conditions sufficient to remove fatty acids or corn oils contained therein;
 - c) selectively extracting pigments and flavor compounds from the destarched and alkaline treated gluten of step (b) with aqueous alcohol having an alcohol concentration that does not substantially extract zein, zein bodies or glutelins to obtain deflavored and decolored glutelins.
24. An essentially decolored and deflavored gluten purified by the method of Claim 23.
25. A food product containing deflavored and decolored gluten produced by the method of Claim 23.

AMENDED CLAIMS

[received by the International Bureau on 14 April 1993 (14.04.93);
original claims 1,4-6,11-14,16-18 and 20 amended; other
claims unchanged (3 pages)]

1. A method of producing decolored and deflavored zein or zein bodies and glutelins from corn gluten comprising removing color and flavor impurities from the gluten by treating corn gluten with alkali to remove fatty acids and corn oils contained therein and then with an aqueous alcoholic solvent having a concentration which does not substantially extract zein, zein bodies and glutelins therefrom, thereafter changing the concentration of the alcoholic solvent to a concentration sufficient to separate the gluten into decolored and deflavored zein or zein bodies and glutelin fractions.
2. The method of Claim 1, wherein the alcohol is a food grade alcohol.
3. The method of Claim 2, wherein the alcohol is ethanol, isopropanol or combinations thereof.
4. The method of Claim 1, wherein the alcohol concentration is from about 90 to about 100% by volume or from about 10 to 40% by volume alcohol and is then changed to a concentration of from about 40 to about 90% by volume alcohol.
5. An edible, essentially decolored and deflavored zein that is white produced by the method of Claim 1.
6. Essentially decolored and deflavored zein bodies that are edible produced by the method of Claim 1.

9. The method of Claim 7, wherein step (b) is performed at a pH of from about 8 to about 9 and a temperature of from about 60 to about 80°C.
10. The method of Claim 7, wherein the alkaline buffer is selected from aqueous solutions of sodium hydroxide, calcium hydroxide, ammonium hydroxide, sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate and mixtures thereof.
11. The method of Claim 7, wherein step (c) is performed with ethanol or isopropanol having an alcohol concentration of from about 90 to about 100% by volume or from about 10 to about 40% by volume.
12. The method of Claim 7, wherein step (d) is performed using an ethanol concentration of from about 40 to from about 90% by volume alcohol or from about 0-40% by volume alcohol.
13. The method of Claim 7, wherein step (e) is performed by precipitating the purified zein in water or from about 10 to about 40% by volume alcohol, removing the zein by filtration therefrom and then drying the product.
14. The method of Claim 7, further comprising washing the product of step (b) with water to remove residual water soluble products from the gluten.
15. The method of Claim 7, further comprising the step of treating the extracted zein with activated carbon or ion exchange resin to remove impurities in the alcohol extracted zein.

16. An edible, essentially decolored and deflavored zein that is white purified by the method of Claim 7.
17. An edible, continuous film cast from an aqueous solution containing decolored and deflavored zein purified by the method of Claim 7.
18. An edible, essentially decolored and deflavored zein body fraction purified by the method of Claim 7.
19. The method of Claim 7, further comprising recovering the glutelin fraction to thereby yield decolored and deflavored glutelins.
20. An ethanol solution of edible, deflavored and decolored zein, purified by the method of Claim 7, which is suitable for use as an edible barrier coating.
21. An essentially decolored and deflavored zein that is suitable for use in a food or pharmaceutical application.
22. A food product containing decolored and deflavored zein, zein bodies, glutelins and combinations thereof.
23. A method of producing deflavored and decolored corn glutens, comprising the steps of:
 - a) destarching the corn gluten with a carbohydrate hydrolyzing enzyme under agitating conditions, sufficient to hydrolyze and remove starch contained therein;




FIG. 1

2/4




FIG. 2

3/4





FIG. 3

4/4

FIG. 4



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US92/11134

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) : A23J 1/12, 3/14; C07K 3/02, 3/12, 3/28, 15/10

US CL : 426/656; 530/373, 374, 424

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 426/656; 530/373, 374, 424

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<u>X</u> Y	US, A, 2,133,591 (Swallen et al.) 18 October 1938, page 1, column 1, lines 48-54; page 2, column 1, lines 46-69, and page 2, column 2, lines 19-23.	<u>5,6,16,18, 20-22</u> 17
X	US, A, 3,802,896 (Westall et al.) 09 April 1974, column 1, lines 13-26, column 4, line 58 - column 5, line 14, and Example III.	5, 6, 16-18, 20
Y	US, A, 3,880,665 (Himmelmann) 29 April 1975, column 14, lines 13-17.	17
X	US, A, 4,018,936 (Garbutt et al.) 19 April 1977, column 1, lines 36-37, and Examples 7 and 12.	24, 25
X	US, A, 4,716,218 (Chen et al.) 29 December 1987, column 2, lines 17-30, and column 3, lines 8-19.	1-6,16,18, 20-22,24,25
A	US, A, 2,479,519 (Schopmeyer) 16 August 1949, column 3, lines 13-20.	1-25



Further documents are listed in the continuation of Box C.



See patent family annex.

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be part of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier document published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"Z"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

25 January 1993

Date of mailing of the international search report

22 FEB 1993

Name and mailing address of the ISA/
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Authorized officer

JEFFREY E. RUSSEL

Facsimile No. NOT APPLICABLE

Telephone No. (703) 308-0196